



CONTINUOUS RESIDUAL ACTIVITY
UP TO 6 MONTHS*

Sporicidin®

BRAND

DISINFECTANT TOWELETTES

An effective disinfectant and active deodorant for hospitals, clinics, medical, dental and veterinary offices, farm premises, laboratories, industrial clean rooms, homes, nursing homes, ambulances, hotels, restaurants, schools, airplanes, trains, boats, autos, buses, health spas and toilets.

Active Ingredients	
Phenol	1.56%
Sodium Phenate	0.06%
Inert Ingredients	98.38%
Total	100.00%

Keep Out of Reach of Children

CAUTION

Net contents: 85 Towelettes 9.5" x 12" saturated with disinfectant; 2 lbs. 7 oz.

- BACTERICIDAL
- FUNGICIDAL
- VIRUCIDAL **
- TUBERCULOCIDAL
- PREVENTS MOLD AND MILDEW
- ELIMINATES ODORS*

EPA REG. NO. 8383-7
EPA Est. Nos. 34346-NE-001,
87937-SC-001, 71979-SC-001
See batch code for actual Est. No.
Part No.: JUM-8506

For viewing purposes only

For viewing purposes only

AN EFFECTIVE CLEANER, DISINFECTANT AND ACTIVE DEODORIZER FOR INSTRUMENTS, AND EQUIPMENT ON PRE-CLEANED, HARD, NON-POROUS SURFACES in health/hospital treatment and patient rooms, operating rooms, schools, ambulances, hotels, restaurants, boats, planes, buses, and the home. Recommended for medical and dental equipment, beds, surgical carts, counter tops, mannequins, telephones, keyboards, furniture, wheelchairs, walkers, sinks, floors, walls, light switches, linen hampers, shower stalls, toilet seats, trash containers and damp and musty areas, including animal areas. Self-sanitizes against odor-causing bacteria and fungi. Reduces the hazards of cross-contamination.

PASSES AOAC EFFICACY STANDARDS FOR DISINFECTION IN THE PRESENCE OF 5% ORGANIC SOIL. FOR INSTITUTIONAL, HOSPITAL AND FARM PREMISE USE.

Virucidal:** Avian Influenza A (H9N2), Influenza A₂ (Japan 305/57 Asian strain), Cytomegalovirus, Coronavirus, Polio Type 1 virus, Vaccinia, Canine parvovirus, Herpes simplex types 1/F and 2/G (oral, ocular and genital), HIV-1 (AIDS Virus). (See Sporidicin Bulletin NO. 303).

Bactericidal: Staphylococcus aureus, Salmonella enterica, Pseudomonas aeruginosa, Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecium (VRE), E. coli, Streptococcus pyogenes, Streptococcus salivarius, Mycobacterium tuberculosis.

Fungicidal: Chaetomium globosum, Aspergillus niger, Penicillium variable, Trichophyton mentagrophytes (Athlete's Foot fungus).

ACTIVE DEODORIZER:** Provides continuous residual bacteriostatic activity against odor-causing bacteria for up to 6 months, if not removed from surfaces including plastics, latex, vinyl, glass, treated wood, metal, glazed porcelain, glazed tile and paint.

SPORICIDIN DISINFECTANT TOWELETTES are germicidal when used as directed for disinfection but differ in composition from other Sporidicin products.

DIRECTIONS FOR USE: It is a violation of U.S. Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT DILUTE OR MIX WITH OTHER CHEMICALS

PRECLEAN surfaces using Sporidicin Disinfectant Solution to remove soil and filth. Wipe dry with paper towel, cloth or sponge.

TO DISINFECT AND DEODORIZE items which cannot be immersed, such as electrical panels, cash registers, door knobs, handles, seats, dashboards and other "high touch" surfaces in and around farm equipment, animal and poultry housing, barns, kennels, breeding pens, hatcheries, trucks and vehicles: Thoroughly wet pre-cleaned surface with Sporidicin Disinfectant Towelettes and allow surface to remain wet for 10 minutes. Use as many towelettes as necessary for the treated area to remain wet for ten minutes at room temperature (20°C/68°F or above) to kill listed organisms.

TO KILL, CLEAN AND PREVENT MOLD AND MILDEW: Use procedure stated above. Repeat application if new growth appears.

TO SPOT CLEAN, DEODORIZE AND REMOVE DEBRIS FROM CARPETS AND FABRICS: Blot or scrub area to be treated with Sporidicin Disinfectant Towelette to remove soiling. Wipe dry with clean cloth or towel.

FOR FOOD PROCESSING PLANTS: Follow pre-cleaning and disinfecting instructions outlined above to disinfect hard, non-porous surfaces and materials in food handling areas, including poultry and meat packing facilities, slaughter houses, etc. All surfaces and materials that come in contact with food must be washed with soap or detergent and rinsed with potable water prior to use.

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS.

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: If in eyes, remove contact lenses. Hold eye open and slowly rinse with water. If irritation persists for more than 5 minutes continue rinsing and consult a physician or poison control center.

STORAGE AND DISPOSAL: Store at room temperature. Nonrefillable container. Do not reuse or refill this container. Discard empty container in trash or offer for recycling if available.

FOR HOSPITAL DISINFECTION: This product is not to be used as a terminal sterilant, high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Made in U.S.A.
Manufactured for:
Contract Corp. / Sporidicin Division
Columbia, SC 29303
1-800-424-3733 www.sporidicin.com